

Ordering temperatures of Ising Spin Glasses

I. A. Campbell¹

¹*Laboratoire des Verres, Université Montpellier II, 34095 Montpellier, France*

(Dated: February 2, 2008)

Exploiting an approach due to Singh and Fisher I show that in the high dimension limit the ordering temperature of near neighbour Ising Spin Glasses drops linearly with the kurtosis of the interaction distribution, in excellent agreement with accurate high temperature series data of Daboul, Chang and Aharony. At lower dimensions the linear relation no longer applies strictly but the kurtosis can still be taken to be an appropriate parameter for ranking different systems. I also compare the series estimates with simulation and Migdal-Kadanoff estimates where these are available.

PACS numbers: 75.50.Lk, 75.40.Mg, 05.50.+q

In order to establish precise values of critical exponents at a continuous transition, it is essential to first obtain accurate and reliable values for the critical temperature T_c . The ordering temperatures T_c of Ising spin glasses (ISGs) are not expected to be universal but to depend on the form of the interaction distribution. In ISGs with random near neighbour interactions of zero mean $\langle J_{ij} \rangle = 0$ the interaction distributions are normalized according to the standard convention $\langle J_{ij}^2 \rangle = 1$ (where $\langle \dots \rangle$ represents an average over the interaction distribution J_{ij} between pairs of spins). It was suggested on purely empirical grounds¹ that T_c values should be ranked from highest to lowest according to the value of the kurtosis of the interaction distribution,

$$K = \langle J_{ij}^4 \rangle / \langle J_{ij}^2 \rangle^2 \quad (1)$$

(An alternative definition of kurtosis, sometimes referred to as the excess kurtosis, corresponds to $K - 3$ which leads to a zero value for the Gaussian distribution. For present purposes it turns out that the the definition in Eq. (1) is more convenient.) Extensive high temperature series calculations by Daboul, Chang and Aharony provide accurate values of T_c in dimensions d from 8 to 4 which are consistent with this phenomenological rule in each dimension².

Exploiting an approach due to Singh and Fisher³ I show rigorously that in the high dimension limit the ordering temperature of near neighbour ISGs drops linearly with the kurtosis of the interaction distribution, and find excellent quantitative agreement between the Singh-Fisher predictions and the data of Daboul *et al*² in dimensions 7 and 8. At lower dimensions the linear relation no longer applies strictly but the trend remains and the kurtosis can still be taken to be an appropriate parameter for ranking different systems.

Singh and Fisher³ used a $1/d$ expansion technique (whose principle largely pre-dates RGT⁴) to evaluate T_c in ISGs as a function of dimension and of interaction distribution. On this approach and for the near neighbour ISG with bimodal $\pm J$ interactions, they define

$$\omega_c(\pm J) = \tanh^2[J/T_c(\pm J)] \quad (2)$$

and find that

$$[\omega_c]^{-0.5} = (\sigma)^{0.5} [1 - (7/2)/\sigma^2 - (21/2)/\sigma^3 - \dots] \quad (3)$$

where $\sigma = (2d - 1)$. They summed this series, truncating appropriately, to estimate the sequence of $T_c(\pm J)$ values for the bimodal $\pm J$ interaction in different dimensions. They obtained satisfactory agreement with the high temperature series values which had just been evaluated for this interaction^{5,6,7}. More recent high temperature series work^{2,8} gives values consistent with the earlier series estimates for the bimodal distribution.

For present purposes the important step of Singh and Fisher was the next one : they state that for an arbitrary interaction distribution that we will label X , the mean over the interaction distribution $J_{ij}(X)$ of

$$\omega_c = \langle \tanh^2[J_{ij}(X)/T_c(X)] \rangle \quad (4)$$

is the same for all distributions to order $1/\sigma^2$. This is an implicit equation for $T_c(X)$ once ω_c is known. The leading deviation was given as

$$[\omega_c(X) - \omega_c(\pm J)] = \omega_c(\pm J) [1 - 3(\mu_2 - 1)/\sigma^3] \dots \quad (5)$$

where

$$\mu_n = \langle \tanh^{2n}[J_{ij}/T_c(X)] \rangle / \langle \tanh^2[J_{ij}/T_c(X)] \rangle^n \quad (6)$$

It turns out that the effect of this deviation term is numerically rather weak in all dimensions.

If d is high enough, T_c values will also be high (in $d = 8$ $T_c \sim 3.6$ in the examples discussed below) so the whole range of $[J_{ij}/T_c(X)]$ values can be taken to be small and we can evaluate $\tanh^2(x)$ for small arguments only. Now

$$\tanh^2(x) = x^2 - (2/3)x^4 + (17/45)x^6 \dots \quad (7)$$

so keeping only the two leading terms, Eq. (4) becomes

$$\omega_c = \langle J_{ij}^2(X) \rangle / T_c(X)^2 - (2/3) \langle J_{ij}^4(X) \rangle / T_c^4 \quad (8)$$

As all ISG interaction distributions are normalized such that $\langle J_{ij}^2(X) \rangle = 1$ and in the high d limit ω_c is independent of distribution,

$$\omega_c = 1/T_c(X)^2 - (2/3)K(X)/(\omega_c T_c(X)^4) \quad (9)$$




FIG. 1: (Colour on line) T_c estimates for near neighbour interaction ISGs in dimension 8 from high temperature series data, Daboul *et al*² (black squares), the Singh-Fisher³ estimates (see text) (green triangles), and the leading approximation Eq. (10), straight line. The points for successive K refer to ISGs with bimodal, double triangle, uniform and Gaussian distributions from left to right (see¹ for the definitions of the interaction distributions with the standard normalizations)

with $K(X)$ the kurtosis of the distribution X . To the same level of approximation the equation can finally be rewritten

$$T_c(X, d) = (1/\omega_c(d))^{0.5} [1 - K(X)\omega_c(d)/3] \quad (10)$$

This should be exact in the limit of high d and small to moderate kurtosis. The critical temperature should then vary linearly with the kurtosis of the distribution $K(X)$, with a slope which is given by $[\omega_c(d)]^{0.5}/3$. $\omega_c(d)$ can be fixed if one $T_c(X)$ (generally the bimodal value) is known.

I calibrate the single free parameter $\omega_c(d)$ by using the Daboul *et al* value for $T_c(\pm J)$ in each dimension (the Singh-Fisher $1/\sigma$ series $T_c(\pm J)$ values are in fact very close to the Daboul *et al* values), except in dimension 4 where the error bar on the Daboul *et al* value is high and I calibrate with the best simulation value. The ω_c values used are 0.069877, 0.08251, 0.1395 and 0.2208 in dimensions 8, 7, 5 and 4 respectively. The distributions studied by Daboul *et al* were the bimodal, double triangle, uniform and Gaussian with $K = 1, 4/3, 9/5$ and 3 respectively. In simulations¹ the Laplacian (or decreasing exponential) distribution which has $K = 6$ was also studied.

In Figures 1 and 2 I show the high temperature series T_c values² plotted against kurtosis for dimensions $d = 8$ and $d = 7$. The leading approximation, Eq. (10), gives the straight line which lies very close to the Daboul *et al* data points for the different distributions. Using the full Singh-Fisher expressions (Eqs. (4), (5), and (6)), evaluated by numerical integration without appealing to the




FIG. 2: (Colour on line) T_c estimates for near neighbour interaction ISGs in dimension 7 from high temperature series data, Daboul *et al*² (black squares), the Singh-Fisher³ estimates (see text) (green triangles), and the leading approximation Eq. (10), straight line. The points refer to ISGs with bimodal, double triangle, uniform and Gaussian distributions from left to right.

small argument approximation gives calculated points which are in almost perfect agreement with the Daboul *et al* data in these dimensions. The weak deviation from the straight line for $K = 3$ (the Gaussian distribution) is due to the error introduced by truncating Eq. (7) in the wings of the Gaussian.

By dimension 5, figure 3, the first order approximation has become much poorer as could be expected *a priori* because the small argument development is no longer valid in this lower dimension, where the T_c values are lower. There are weak deviations of the Singh-Fisher estimates (without the small argument approximation) from the high temperature series data points, meaning that Eq. (5) is no longer accurately obeyed, either because higher order terms give non-negligible contributions or more fundamentally because the $1/\sigma$ expansion becomes inaccurate at lower dimensions. As pointed out by Singh and Fisher their expansion "is at best asymptotic and optimum estimates come by truncating after the smallest term"³, a procedure which introduces uncertainties at lower dimensions. Nevertheless the general trend of decreasing T_c with increasing kurtosis is still clearly present.

Finally for $d = 4$ the small argument approximation to the Singh-Fisher expressions is certainly inappropriate but one can again evaluate the full Singh-Fisher ordering temperature estimates through numerical integration. In this dimension ordering temperatures have also been estimated independently by simulations, using various complementary methods. These include a technique relying on consistency between dynamic and static critical parameters¹, domain wall analysis¹⁰, and Binder param-




FIG. 3: (Colour on line) T_c estimates for near neighbour interaction ISGs in dimension 5 from high temperature series data, Daboul et al² (black squares), the Singh-Fisher³ estimates (see text) (green triangles), and the leading approximation Eq. (10), straight line. The points refer to ISGs with bimodal, double triangle, uniform and Gaussian distributions from left to right.

eter measurements^{11,12,13,14}. In the two cases (bimodal and Gaussian interaction distributions) where simulation results using different methods exist, agreement between the various simulation estimates is excellent, see Table I. In addition, the T_c values have been studied in $d = 4$ (and $d = 3$) using a Migdal-Kadanoff (MK) approach⁹ where the optimum size b^* of the MK renormalization cell is chosen by interpolating the number of segments b in an MK branch until $T_c(MK)$ is equal to T_c for the $\pm J$ distribution. $T_c(MK)$ values for the other distributions are then estimated with this same b^* . For $d = 4$ the optimum interpolated value for b was found to be $b^* = 2.49^9$. The different T_c estimates are given in Table 1 and are shown in Figure 4 where the same general trend once more appears. It can be seen that while the Singh-Fisher values and the MK values lie close to but not precisely on the series and simulation points, the agreement between the simulation estimates and the series values is excellent. As the simulation calculations are necessarily carried out on samples of moderate size (a maximum of $L = 12$ up to now in dimension 4), *a priori* there might have been a lingering suspicion that the T_c estimates from simulations could somehow be intrinsically in error because of corrections to finite size scaling that had not been properly taken into account. Because the series estimates are not subject to this type of error, the agreement between the two sets of quite independent T_c values lays this suspicion to rest. One can have confidence that at least in dimension $d = 4$ the simulation estimates of $T_c(X)$ (and in particular those using the static/dynamic consistency technique¹ which are the most complete) are intrinsically




FIG. 4: (Colour on line) T_c estimates for near neighbour interaction ISGs in dimension 4 from high temperature series data, Daboul et al² (black squares), the Singh-Fisher³ estimates (see text) (green triangles), simulation estimates¹ (red circles), and MK estimates with $b^* = 2.49^9$ (blue inverted triangles). The points refer to ISGs with bimodal, double triangle, uniform, Gaussian and Laplacian distributions from left to right.

TABLE I: Comparison of estimates of T_c in dimension 4

distribution	method	reference	T_c
bimodal($K = 1$)	series	⁸	2.02(3)
bimodal	series	²	1.96(7)
bimodal	simulation	¹⁴	2.03(3)
bimodal	simulation	¹⁰	2.00(4)
bimodal	simulation	¹¹	2.00(1)
bimodal	simulation	¹	1.99(1)
triangles($K = 4/3$)	series	²	1.95(1)
uniform($K = 9/5$)	series	²	1.90(2)
uniform	simulation	¹	1.91(1)
uniform	MK	⁹	1.87(2)
Gaussian($K = 3$)	series	²	1.78(1)
Gaussian	simulation	¹³	1.80(2)
Gaussian	simulation	¹²	1.80(2)
Gaussian	simulation	¹	1.76(2)
Gaussian	MK	⁹	1.75(2)
Laplacian($K = 6$)	simulation	¹	1.52(1)
Laplacian	MK	⁹	1.54(2)

correct and are not polluted by uncontrolled corrections to finite size scaling.

Empirically, Table 1 shows that the MK T_c values are a good indication of the true T_c values if the appropriate choice of b^* has been made by calibrating b using the T_c for one particular distribution (see the discussion in⁹). In dimension 3 where the series method would require a very long set of approximants, and where it turns out

that in simulations corrections to finite size scaling are much stronger than in dimension 4, it would appear that the ratios of T_c values for different distributions obtained through the MK approach^{9,15} should be rather reliable and a very helpful indication for controlling simulation measurements.

We can conclude that the key to understanding how T_c values vary from distribution to distribution in ISGs with near neighbour interactions is the Singh-Fisher statement that $\omega_c(d)$ (Eq. (4)) is almost distribution independent for fixed dimension d . The linear dependence of $T_c(X)$ on the kurtosis $K(X)$ with a slope $[\omega_c(d)]^{0.5}/3$ becomes exact in the high dimension limit and is still a close approximation in high but finite dimensions ($d = 8$ and 7). Even at moderate d ($d = 5$ and 4), the general trend still holds so the ranking of distributions by their kurtosis remains physically appropriate in agreement with the

initial phenomenological suggestion¹. (It is nevertheless conceivable that for low dimensions differences between distributions could lead to situations such that two distributions with identical kurtoses would have marginally different T_c values. For the moment the number of different distributions that have been studied is very limited so this possibility has never been put to the test). In dimension $d = 4$ which is the only dimension where appropriate data is available for both simulations and series measurements, simulation estimates of ordering temperatures are in excellent agreement with the series values which confirms that the simulation estimates, in particular those based on the consistency between dynamic and static critical parameters¹, are accurate and reliable.

Acknowledgements : I would like to thank A. Aharony, D. Daboul, R.R.P. Singh and H.G. Katzgraber for their helpful comments.

-
- ¹ L.W. Bernardi and I.A. Campbell, Phys.Rev.B **56**, 5271 (1997)
 - ² D. Daboul, I. Chang, and A. Aharony, Europhys. J. B **41**, 231 (2004)
 - ³ R.R.P. Singh and M.E. Fisher, J. App. Phys. **63**, 3994 (1988)
 - ⁴ M.E. Fisher and D.S. Gaunt, Phys. Rev. A **133**, 224 (1964)
 - ⁵ R.R.P. Singh and S. Chakravarty, Phys. Rev. Lett. **57**, 245 (1986)
 - ⁶ R.R.P. Singh and S. Chakravarty, Phys. Rev. B **36**, 546 (1987)
 - ⁷ R.R.P. Singh and S. Chakravarty, Phys. Rev. B **36**, 559 (1987)

- ⁸ L. Klein, J. Adler, A. Aharony, A.B. Harris and Y. Meir, Phys. Rev. B **43**, 11249 (1991)
- ⁹ S. Prakash and I.A. Campbell, Physica A **235**, 507 (1997)
- ¹⁰ K. Hukushima, Phys. Rev. E **60**, 3606 (1999)
- ¹¹ A.P. Young, unpublished
- ¹² M. Ney-Nefle, Phys. Rev. B **57**, 492 (1998)
- ¹³ G. Parisi, F. Ricci-Tersenghi, J. J. Ruiz-Lorenzo, J. Phys. A **29**, 7943 (1996)
- ¹⁴ E. Marinari and F. Zuliani, J. Phys. A **32**, 7447 (1999)
- ¹⁵ E. Nogueira, S. Coutino, F.D. Nobre and E.M.F. Curado, Physica A **271**, 125 (1999)